

Vertex integrals in heavy-particle theories

Antonio O. Bouzas

Departamento de Física Aplicada, Cinvestav-IPN, Apdo. Postal 73 “Cordemex”, Mérida 97310, Yucatán, México

Abstract. We give the results of complete analytical computations of two- and three-point loop integrals occurring in heavy particle theories, with and without velocity change, for arbitrary values of external momenta and masses.

INTRODUCTION

In this talk we consider a class of one-loop integrals occurring in heavy-particle theories [1], with arbitrary real values for the external masses and residual momenta. We give the results of complete analytical computations of three-point loop integrals with and without velocity change, and two-point loop integrals. The details of the calculations are given in [2], and in a forthcoming paper.

LOOP INTEGRALS

The loop integrals we consider are of the form,

$$\begin{aligned}
 j_2^{\alpha_1 \dots \alpha_n} &= \frac{i\mu^{4-d}}{(2\pi)^d} \int d^d \ell \frac{\ell^{\alpha_1} \dots \ell^{\alpha_n}}{(2v \cdot (\ell + k) - \delta M + i\epsilon)(\ell^2 - m^2 + i\epsilon)} \\
 j_3^{\alpha_1 \dots \alpha_n} &= \frac{i\mu^{4-d}}{(2\pi)^d} \int d^d \ell \frac{\ell^{\alpha_1} \dots \ell^{\alpha_n}}{(2v_1 \cdot (\ell + k_1) - \delta M_1 + i\epsilon)(2v_2 \cdot (\ell + k_2) - \delta M_2 + i\epsilon)} \\
 &\quad \times \frac{1}{(\ell^2 - m^2 + i\epsilon)} \\
 \mathcal{H}_3^{\alpha_1 \dots \alpha_n} &= \frac{i\mu^{4-d}}{(2\pi)^d} \int d^d \ell \frac{\ell^{\alpha_1} \dots \ell^{\alpha_n}}{(2v \cdot \ell - \delta M + i\epsilon)((\ell - k_1)^2 - m^2 + i\epsilon)} \\
 &\quad \times \frac{1}{((\ell - k_2)^2 - m'^2 + i\epsilon)}
 \end{aligned} \tag{1}$$

Here v_i^μ , $i = 1, 2$, are the velocities of the external heavy legs, k_i^μ their residual momenta, and δM_i their mass splittings relative to the common heavy mass of the corresponding heavy quark symmetry multiplet. m and m' are the masses of the light particles within the loops, which in chiral theories are light pseudoscalar mesons. These integrals are defined in $d = 4 - \epsilon$ dimensions, μ being the mass scale of dimensional regularization.

Their degrees of divergence are $n + d - 3$ for $j_2^{\alpha_1 \cdots \alpha_n}$, $n + d - 4$ for $j_3^{\alpha_1 \cdots \alpha_n}$ and $n + d - 5$ for $\mathcal{H}_3^{\alpha_1 \cdots \alpha_n}$. The factor of 2 in front of v_i^μ corresponds to our normalization of the heavy-particle propagators.

Our method of calculation [2] is to obtain the integrals (1) as large-mass limits of ordinary loop integrals. We closely follow the approach of [3] for the computation of scalar integrals, which are greatly simplified in the large-mass limit, and the method of [4] to express tensor integrals in terms of scalar ones.

TWO-POINT INTEGRALS

Two-point integrals with one heavy propagator have been given in [2, 5, 6]. The scalar two-point integral j_2 is a function of m and $\Delta = \delta M - 2v \cdot k$. We write it in terms of $\xi = \Delta/(2m)$,

$$j_2(\Delta, m) = \frac{\Delta}{32\pi^2} \left(\frac{2}{\varepsilon} + \log\left(\frac{\bar{\mu}^2}{m^2}\right) + 2 \right) + \frac{m}{16\pi^2} \mathcal{F}(\xi)$$

with $\mathcal{F}(x) = \sqrt{x^2 - 1 + i\varepsilon} \left[\log\left(x - \sqrt{x^2 - 1 + i\varepsilon}\right) - \log\left(x + \sqrt{x^2 - 1 + i\varepsilon}\right) \right]$. The coefficient of the dimensional regularization pole vanishes when $\Delta = 0$. This is due to the fact that the real part of the integrand in j_2 is parity-odd when $\Delta = 0$.

The vector two-point integral $j_2^\mu(v^\alpha, \Delta, m)$ is given in terms of only one form factor, $j_2^\mu(v^\alpha, \Delta, m) = F(\Delta, m)v^\mu$, with $F(\Delta, m) = v_\mu j_2^\mu(v^\alpha, \Delta, m) = 1/2A_0(m) + \Delta/2j_2(\Delta, m)$, where A_0 is the standard one-point scalar integral (see the appendix of [2]). The second-rank tensor integral is computed analogously, explicit results being given in [2].

THREE-POINT INTEGRALS WITH VELOCITY CHANGE

The scalar three-point integral with velocity change, $j_3 = j_3(v_1 \cdot v_2, \Delta_1, \Delta_2, m)$, where $\Delta_j = \delta M_j - 2v_j \cdot k_j$, can be expressed in terms of four dilogarithms [2],

$$j_3 = \frac{1}{64\pi^2} \frac{1 - \Omega^2}{\Omega} \left\{ \frac{2}{\varepsilon} \log(\alpha) + \log^2(\alpha) + \sum_{k, \sigma} (-1)^{k+1} \left[\frac{1}{2} \log^2\left(\frac{-z_{k\sigma}}{\bar{\mu}}\right) + \text{Li}_2\left(\frac{-y_0}{z_{k\sigma}}\right) \right] \right\}.$$

The notation is as follows, $\omega = v_1 \cdot v_2$, $\Omega = \sqrt{(\omega - 1)/(\omega + 1)}$, $\alpha = \omega + \sqrt{\omega^2 - 1} = (1 + \Omega)/(1 - \Omega)$ and $\bar{\mu}$ is the mass unit in the $\overline{\text{MS}}$ scheme. The sum extends over $k = 1, 2$ and $\sigma = \pm$, with

$$\begin{aligned} y_0 &= -\frac{1 + \Omega}{2\Omega} \left(\frac{1 + \Omega}{2} \Delta_1 - \frac{1 - \Omega}{2} \Delta_2 \right) \\ z_{1\pm} &= \frac{1}{2} \left(\frac{1 + \Omega^2}{2\Omega} \Delta_1 - \frac{1 - \Omega^2}{2\Omega} \Delta_2 \pm \sqrt{\Delta_1^2 - 4m^2 + i\varepsilon} \right) \\ z_{2\pm} &= \frac{\alpha}{2} \left(\frac{1 - \Omega^2}{2\Omega} \Delta_1 - \frac{1 + \Omega^2}{2\Omega} \Delta_2 \pm \sqrt{\Delta_2^2 - 4m^2 + i\varepsilon} \right). \end{aligned}$$

This expression for $\mathcal{J}_3$ is equivalent to the result given in eq. (30) of [2], though it has been written in a more compact form by means of the identity

$$\frac{1}{2}\log^2(z) - \log(z)\log(-z) = -\frac{\pi^2}{2} - \frac{1}{2}\log^2(-z),$$

valid on the first Riemann sheet of the logarithm, and the identity (A.2) of [3] for the dilogarithm.

In order to compute the vector integral $\mathcal{J}_3^\mu$ we define two sets of form factors as $\mathcal{J}_3^\mu = I_1 v_1^\mu + I_2 v_2^\mu$ and $F_{1,2} = v_{1,2} \cdot \mathcal{J}_3$. These form factors are given by,

$$I_1 = \frac{1 - \Omega^2}{4\Omega^2} [-(1 - \Omega^2)F_1 + (1 + \Omega^2)F_2], \quad I_2 = \frac{1 - \Omega^2}{4\Omega^2} [(1 + \Omega^2)F_1 - (1 - \Omega^2)F_2],$$

with $F_{1,2} = 1/2 \mathcal{J}_2(\Delta_{2,1}, m) + \Delta_{1,2}/2 \mathcal{J}_3(v_1 \cdot v_2, \Delta_1, \Delta_2, m)$. These equations give an explicit expression for $\mathcal{J}_3^\mu$. For the sake of brevity, we omit here the results for the tensor integral $\mathcal{J}_3^{\mu\nu}$, which can be found in [2].

THREE-POINT INTEGRALS WITH ONE HEAVY PROPAGATOR

The scalar three-point integral $\mathcal{H}_3 = \mathcal{H}_3(v \cdot q, q^2, \Delta, m, m')$, with $q = (k_2 - k_1)/2$ and $\Delta = \delta M - v \cdot (k_1 + k_2)$, can be expressed in terms of eight dilogarithms as,

$$\begin{aligned} \mathcal{H}_3 &= \frac{1}{(4\pi)^2} \frac{1}{4|\mathbf{q}|} \sum_{j=1,2} (\mathcal{F}_1(y_j) + \mathcal{F}_2(x_j) - \mathcal{F}_3(z_j)) \\ \mathcal{F}_1(x) &= \text{Li}_2\left(\frac{z_0 - 4|\mathbf{q}|\alpha}{z_0 - 4|\mathbf{q}|x}\right) - \text{Li}_2\left(\frac{z_0}{z_0 - 4|\mathbf{q}|x}\right) \\ \mathcal{F}_2(x) &= -\text{Li}_2\left(\frac{z_0 - 4|\mathbf{q}|\alpha}{z_0 - 4|\mathbf{q}|x}\right) - \frac{1}{2}\log^2\left(\frac{z_0 - 4|\mathbf{q}|x}{\mu^2}\right) \\ \mathcal{F}_3(x) &\equiv \mathcal{F}_1(x) + \mathcal{F}_2(x) = -\text{Li}_2\left(\frac{z_0}{z_0 - 4|\mathbf{q}|x}\right) - \frac{1}{2}\log^2\left(\frac{z_0 - 4|\mathbf{q}|x}{\mu^2}\right). \end{aligned}$$

The quantities entering these equations are $|\mathbf{q}| = \sqrt{(v \cdot q)^2 - q^2}$ ($|\mathbf{q}|$ is assumed to be real), $\alpha \equiv \alpha_+$ with $\alpha_{\pm} = 2(v \cdot q \pm |\mathbf{q}|)$, $z_0 = -(m'^2 - m^2) - \alpha_+(\Delta - 2|\mathbf{q}|)$, and,

$$\begin{aligned} x_{1,2} &= v \cdot k_2 + 2|\mathbf{q}| - \frac{\delta M}{2} \pm \sqrt{\left(v \cdot k_1 - \frac{\delta M}{2}\right)^2 - m^2 + i\epsilon} \\ y_{1,2} &= \frac{1}{2\alpha_-} \left(4q^2 + m'^2 - m^2 \pm \sqrt{(4q^2 - (m' + m)^2)(4q^2 - (m' - m)^2) + i\epsilon\sigma}\right) \\ z_{1,2} &= v \cdot k_2 - \frac{\delta M}{2} \pm \sqrt{\left(v \cdot k_2 - \frac{\delta M}{2}\right)^2 - m'^2 + i\epsilon}, \end{aligned}$$

where in the expression for y_j we denoted $\sigma \equiv \text{sgn}(q^2)$. μ is a positive constant with dimension of mass, analogous to the mass unit in dimensional regularization. It is not difficult to show that $\mathcal{H}_3$ does not depend on μ , it appears there for purely dimensional reasons.

Tensor three-point integrals $\mathcal{H}_3^{\alpha_1 \cdots \alpha_n} = \mathcal{H}_3^{\alpha_1 \cdots \alpha_n}(v, k_1, k_2; \delta M, m, m')$ can be given in terms of integrals with smaller ranks and fewer points, by the well-known method of [4]. We will consider integrals of standard form $H_3^{\alpha_1 \cdots \alpha_n}(v, q; \Delta, m, m') = \mathcal{H}_3^{\alpha_1 \cdots \alpha_n}(v, -q, q; \Delta + v \cdot (k_1 + k_2), m, m')$ in terms of which we can express $\mathcal{H}_3$ as,

$$\mathcal{H}_3^{\alpha_1 \cdots \alpha_n} = H_3^{\alpha_1 \cdots \alpha_n}(v, q; \Delta, m, m') + \sum_{j=1}^n r^{\{\alpha_1 \dots \alpha_j\}} r^{\alpha_{j+1} \cdots \alpha_n} H_3^{\alpha_{j+1} \cdots \alpha_n}(v, q; \Delta, m, m'),$$

where $r^\mu \equiv 1/2(p' + p)^\mu$, and $A^{\{\alpha_1 \alpha_2 \cdots \alpha_s\}} \equiv A^{\alpha_1 \alpha_2 \cdots \alpha_s} + A^{\alpha_2 \cdots \alpha_s \alpha_1} + \dots + A^{\alpha_s \alpha_1 \cdots \alpha_{s-1}}$. Clearly, for the scalar integral we have $\mathcal{H}_3 = H_3$.

For the vector integral we write $H_3^\alpha(v, q; \Delta, m_1, m_2) = V v^\alpha + Q q^\alpha$, with $V = V(v \cdot q, q^2, \Delta, m_1, m_2)$ and similarly Q . If $(v \cdot q)^2 - q^2 = |\mathbf{q}|^2 = 0$, then $q^\alpha \propto v^\alpha$ and we can set $Q = 0$. If $|\mathbf{q}|^2 \neq 0$,

$$|\mathbf{q}|^2 Q = v \cdot q v_\alpha H_3^\alpha - q_\alpha H_3^\alpha, \quad |\mathbf{q}|^2 V = -q^2 v_\alpha H_3^\alpha - v \cdot q q_\alpha H_3^\alpha,$$

with,

$$\begin{aligned} v_\alpha H_3^\alpha &= \frac{1}{2} B_0(4q^2, m_1, m_2) + \frac{\Delta}{2} H_3 \\ q_\alpha H_3^\alpha &= \frac{1}{4} j_2(\Delta - 2v \cdot q, m_2) - \frac{1}{4} j_2(\Delta + 2v \cdot q, m_1) + \frac{m_1^2 - m_2^2}{4} H_3. \end{aligned}$$

We have omitted the arguments $(v, q; \Delta, m_1, m_2)$ of H_3^α on both sides of these equations for brevity. B_0 is the standard scalar two-point integral, as given in the appendix of [2]. Higher-rank tensor integrals can be computed analogously.

The results presented in this section were obtained in collaboration with R. Flores Mendieta. A detailed derivation will be given elsewhere.

ACKNOWLEDGEMENTS

This work has been partially supported by Conacyt and SNI.

REFERENCES

1. See, *e.g.*, A.Manohar, M.Wise, “*Heavy Quark Physics*,” Cambridge Univ. Press, 2000, and references therein.
2. A.Bouzas, Eur.Phys.J. **C12**, (2000), 643
3. G.’t Hooft, M.Veltman, Nucl.Phys. **B153**, (1979), 365.
4. G.Passarino, M.Veltman, Nucl.Phys. **B160**, (1979), 151.
5. D.Broadhurst, A.Grozin, Phys.Lett. **B267**, (1991), 105.
6. I.W.Stewart, Nucl.Phys. **B529**, (1998), 62.